

Metadata

Dataset Name	Northern Hemisphere Extratropical Cyclone Tracks from ERA-5
Dataset General Type	cyclone tracks
Dataset Type	Dataset
Dataset Level	1.1
Program Website	
Keyword Vocabulary	Polar Data Catalogue
Keyword Vocabulary URL	https://www.polardata.ca/pdcinput/public/keywordlibrary
Theme	
Title	Atmosphere
URL	https://canwin-datahub.ad.umanitoba.ca/data/en/group/modelling
Dataset Status	Complete
Maintenance and Update Frequency	As needed
Dataset Last Revision Date	2025-09-23
Dataset DOI	10.34992/ebnw-s681
Metadata Creation Date	2025
Publisher	CanWIN

Dataset Authors

Dataset Authors 1

Name	Crawford, Alex
Type of Name	Personal
Email	alex.crawford@umanitoba.ca
Affiliation	Agriculture and Agri Food Canada
ORCID ID	0000-0003-1561-290X
	ORCID
	http://orcid.org/

Contributors

Contributors 1

Name	Serreze, Mark C
Role	ProjectMember
Email	
Affiliation	National Snow and Ice Data Center, Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder
ORCID ID	0000-0002-3699-302X
	ORCID
	http://orcid.org/

Contributors 2

Name	Sommer, Nathan
Role	ProjectMember
Email	nsommer@wooster.edu
Affiliation	College of Wooster
ORCID ID	ORCID
	ORCID
	http://orcid.org/

Project Data Curator

Alex D Crawford

Project Data Curator email

alex.crawford@umanitoba.ca

Project Data	
Curator Affiliation	Centre for Earth Observation Science - University of Manitoba
Dataset Collection Start Date	1979-01-01
Dataset Collection End Date	2024-12-31
Sample Collection	
Activity Collection Type	
Preferred citation	
Analytical Instrument	
Analytical Method	
License Name	Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International
Licence Type	Open
Embargo Date	
Licence URL	https://spdx.org/licenses
Terms of Access	CanWIN datasets are licensed individually, however most are licensed under the Creative Commons Attribution 4.0 International (CC BY 4.0) Public License. Details for the licence applied can be found using the Licence URL link provided with each dataset. By using data and information provided on this site you accept the terms and conditions of the License. Unless otherwise specified, the license grants the rights to the public to use and share the data and results derived therefrom as long as the proper acknowledgment is given to the data licensor (citation), that any alteration to the data is clearly indicated, and that a link to the original data and the license is made available.
Terms of Use	By accessing this data you agree to [CanWIN's Terms of Use](https://dev.uni-manitoba.links.com.au/data/publication/canwin-data-statement/resource/5b942a87-ef4e-466e-8319-f588844e89c0).
Awards	

Related Resources

Related Resources 1

Related Resource Name	
Resource Code	10.24381/cds.adbb2d47
Identifier Type	
Relationship To This Dataset	
Resource Type	Online Resource
Type	Model
Series Name	

Related Resources 2

Related Resource Name	
Resource Code	
Identifier Type	
Relationship To This Dataset	
Resource Type	Online Resource
Type	
Series Name	

Related Resources 3

Related Resource Name	Mean Pressure at Sea-level from ERA-5
Resource Code	
Identifier Type	DOI
Relationship To This Dataset	IsRequiredBy

Resource Type Online Resource

Type

Series Name

Related Resources 4

Related Resource Name ETOP01 Ice Surface

Resource Code 10.7289/V5C8276M

Identifier Type DOI

Relationship To This Dataset IsReferencedBy

Resource Type Online Resource

Type Dataset

Series Name

Related Resources 5

Related Resource Name CEOS/NSIDC Cyclone Detection and Tracking Algorithm

Resource Code 10.5281/zenodo.4356161

Identifier Type DOI

Relationship To This Dataset IsRequiredBy

Resource Type Online Resource

Type Software

Series Name

Publications

Publications 1

Publication Name Estimating Southern Ocean Storm Positions With Seismic Observations

Identifier Code <https://doi.org/10.1029/2019JC015898>

Identifier Type DOI

Relationship to this dataset IsSupplementedBy

Resource Type Online Resource

Publication Type JournalArticle

Publications 2

Publication Name Sea ice loss and Arctic cyclone activity from 1979 to 2014

Identifier Code <https://doi.org/10.1175/JCLI-D-16-0542.1>

Identifier Type DOI

Relationship to this dataset IsSupplementedBy

Resource Type Online Resource

Publication Type JournalArticle

Publications 3

Publication Name Does the summer Arctic Frontal Zone influence Arctic Ocean cyclone activity?

Identifier Code

Identifier Type DOI

Relationship to this dataset IsSupplementedBy

Resource Type Online Resource

Publication Type

Publications 4

Publication Name Projected Changes in the Arctic Frontal Zone and Summer Arctic Cyclone Activity in the CESM Large Ensemble

Identifier Code <https://doi.org/10.1175/JCLI-D-17-0296.1>

Identifier Type DOI

Relationship to this dataset IsSupplementedBy

Resource Type Online Resource

Publication Type JournalArticle

Publications 5

Publication Name Synoptic Climatology of Rain-on-Snow Events in Alaska

Identifier Code <https://doi.org/10.1175/MWR-D-19-0311.1>

Identifier Type DOI

Relationship to this dataset IsSupplementedBy

Resource Type Online Resource

Publication Type JournalArticle

Publications 6

Publication Name Impacts of synoptic-scale cyclones on Arctic sea-ice concentration: a systematic analysis

Identifier Code <https://doi.org/10.1017/aog.2020.23>

Identifier Type DOI

Relationship to this dataset IsSupplementedBy

Resource Type Online Resource

Publication Type JournalArticle

Publications 7

Publication Name Sensitivity of Northern Hemisphere Cyclone Detection and Tracking Results to Fine Spatial and Temporal Resolution Using ERA5

Identifier Code <https://journals.ametsoc.org/view/journals/mwre/149/8/MWR-D-20-0417.1.xml>

Identifier Type URL

Relationship to this dataset IsCitedBy

Resource Type Online Resource

Publication Type JournalArticle

Publications 8

Publication Name The Influence of the Arctic Frontal Zone on Summer Cyclone Activity Today and in the Future (Doctoral Dissertation)

Identifier Code https://scholar.colorado.edu/concern/graduate_thesis_or_dissertations/6395w720f

Identifier Type URL

Relationship to this dataset IsContinuedBy

Resource Type Online Resource

Publication Type Dissertation

Spatial regions northern-hemisphere

Spatial extent West Bound Longitude	-180.0
Spatial extent East Bound Longitude	180.0
Spatial extent South Bound Latitude	0.0
Spatial extent North Bound Latitude	90.0

Data and Resources

URL	https://zenodo.org/records/7562953
Name	CEOS/NSIDC Extratropical Cyclone Tracking (CNECT) Algorithm
Description	This algorithm has two steps: 1) detection of cyclone centers and areas and 2) tracking of those features. Center detection is based on local minima in sea-level pressure (within a 200 km radius) that have a pressure gradient of at least 7.5 hPa/1000 km. The area of storms and presence of single- and multi-center cyclones are determined using last-closed isobars. Tracking is based on the nearest neighbor to a predicted cyclone propagation location. Cyclone size, intensity, propagation, and interactions (e.g., splitting and merging with other storms) are tabulated at each observation time.
Format	Python
Resource Category	scripts
URL	https://canwin-datahub.ad.umanitoba.ca/data/dataset/4be4d01a-a14b-483f-a1a6-6ead0974fa57/resource/69811381-b58c-4621-b73d-baf1758706f0/download/supplemental-metadata.pdf
Name	Supplemental Metadata
Description	Additional metadata, which includes variable headers, units, and descriptions, as well as an overview of the script applied.
Format	PDF
Resource Category	documents

URL	https://canwin-datahub.ad.umanitoba.ca/data/dataset/4be4d01a-a14b-483f-a1a6-6ead0974fa57/resource/a7ed8d55-63d1-4d7f-aa6f-63fc106e2176/download/cycloneparams.pkl
Name	Cyclone Parameters File
Description	This cyclone parameters file records the input parameters used for the cyclone detection and tracking code to produce the files in this database. It can be opened using pandas in Python via <code>pandas.read_pickle(\$FILEPATH\$)</code> , where <code>\$FILEPATH\$</code> is the path to where this file is stored on your computer.
Format	pkl
Resource Category	supplemental

URL	https://canwin-datahub.ad.umanitoba.ca/data/dataset/4be4d01a-a14b-483f-a1a6-6ead0974fa57/resource/c31c6ec8-2fa8-4d62-8c4f-0ee2d18beedd/download/cnect-nh-era5-1940-1949.zip
Name	Northern Hemisphere Extratropical Cyclone Tracks from ERA5: 1940-1949
Description	Folder containing CSV files that describe extratropical cyclone tracks detected from the ERA5 atmospheric analysis using version 13.2 of the CEOS/NSIDC extratropical cyclone tracking algorithm (as described in Crawford et al., 2021; https://doi.org/10.1175/MWR-D-20-0417.1). Storms are detected with a 3-h temporal resolution and 25-km spatial resolution for the period 1940-1949. All storms whose ending time (cyclolysis) occurs in the same month are grouped into a single CSV file and can be identified by their unique system number (sid). The monthly files are organized within folders for each year. See accompanying metadata file for more details (e.g., descriptions and units for each column). This algorithm explicitly tracks each center of a multi-center cyclones, but in this database, each multi-center cyclone is represented only by its primary center -- i.e., the entire storm system is being represented by the location and area. Only storm systems that meet the following four criteria are included: lifespan of at least 24 hours, track length of at least 1000 km, observed at least once over an elevation less than 500 m, and observed at least once at a distance of at least 500 km from its origin point are included.
Format	ZIP
Resource Category	data

URL	https://canwin-datahub.ad.umanitoba.ca/data/dataset/4be4d01a-a14b-483f-a1a6-6ead0974fa57/resource/d9f63012-7041-48ba-bed2-1bf6e77fa75d/download/cnect-nh-era5-1950-1959.zip
Name	Northern Hemisphere Extratropical Cyclone Tracks from ERA5: 1950-1959
Description	Folder containing CSV files that describe extratropical cyclone tracks detected from the ERA5 atmospheric analysis using version 13.2 of the CEOS/NSIDC extratropical cyclone tracking algorithm (as described in Crawford et al., 2021; https://doi.org/10.1175/MWR-D-20-0417.1). Storms are detected with a 3-h temporal resolution and 25-km spatial resolution for the period 1950-1959. All storms whose ending time (cyclolysis) occurs in the same month are grouped into a single CSV file and can be identified by their unique system number (sid). The monthly files are organized within folders for each year. See accompanying metadata file for more details (e.g., descriptions and units for each column). This algorithm explicitly tracks each center of a multi-center cyclones, but in this database, each multi-center cyclone is represented only by its primary center -- i.e., the entire storm system is being represented by the location and area. Only storm systems that meet the following four criteria are included: lifespan of at least 24 hours, track length of at least 1000 km, observed at least once over an elevation less than 500 m, and observed at least once at a distance of at least 500 km from its origin point are included.

Format	ZIP
Resource Category	data
URL	https://canwin-datahub.ad.umanitoba.ca/data/dataset/4be4d01a-a14b-483f-a1a6-6ead0974fa57/resource/d0abcf90-1a9e-4aa4-b533-19e1b5f54c51/download/cnect-nh-era5-1960-1969.zip
Name	Northern Hemisphere Extratropical Cyclone Tracks from ERA5: 1960-1969
Description	Folder containing CSV files that describe extratropical cyclone tracks detected from the ERA5 atmospheric analysis using version 13.2 of the CEOS/NSIDC extratropical cyclone tracking algorithm (as described in Crawford et al., 2021; https://doi.org/10.1175/MWR-D-20-0417.1). Storms are detected with a 3-h temporal resolution and 25-km spatial resolution for the period 1960-1969. All storms whose ending time (cyclolysis) occurs in the same month are grouped into a single CSV file and can be identified by their unique system number (sid). The monthly files are organized within folders for each year. See accompanying metadata file for more details (e.g., descriptions and units for each column). This algorithm explicitly tracks each center of a multi-center cyclones, but in this database, each multi-center cyclone is represented only by its primary center -- i.e., the entire storm system is being represented by the location and area. Only storm systems that meet the following four criteria are included: lifespan of at least 24 hours, track length of at least 1000 km, observed at least once over an elevation less than 500 m, and observed at least once at a distance of at least 500 km from its origin point are included.
Format	ZIP
Resource Category	data
URL	https://canwin-datahub.ad.umanitoba.ca/data/dataset/4be4d01a-a14b-483f-a1a6-6ead0974fa57/resource/a1928b10-3cc9-41d2-8f59-9bc213373468/download/cnect-nh-era5-1970-1979.zip
Name	Northern Hemisphere Extratropical Cyclone Tracks from ERA5: 1970-1979
Description	Folder containing CSV files that describe extratropical cyclone tracks detected from the ERA5 atmospheric analysis using version 13.2 of the CEOS/NSIDC extratropical cyclone tracking algorithm (as described in Crawford et al., 2021; https://doi.org/10.1175/MWR-D-20-0417.1). Storms are detected with a 3-h temporal resolution and 25-km spatial resolution for the period 1970-1979. All storms whose ending time (cyclolysis) occurs in the same month are grouped into a single CSV file and can be identified by their unique system number (sid). The monthly files are organized within folders for each year. See accompanying metadata file for more details (e.g., descriptions and units for each column). This algorithm explicitly tracks each center of a multi-center cyclones, but in this database, each multi-center cyclone is represented only by its primary center -- i.e., the entire storm system is being represented by the location and area. Only storm systems that meet the following four criteria are included: lifespan of at least 24 hours, track length of at least 1000 km, observed at least once over an elevation less than 500 m, and observed at least once at a distance of at least 500 km from its origin point are included.
Format	ZIP
Resource Category	data
URL	https://canwin-datahub.ad.umanitoba.ca/data/dataset/4be4d01a-a14b-483f-a1a6-6ead0974fa57/resource/6240a835-2ea4-470a-9cc7-7d2ff8276bcb/download/cnect-nh-era5-1980-1989.zip

Name	Northern Hemisphere Extratropical Cyclone Tracks from ERA5: 1980-1989
Description	Folder containing CSV files that describe extratropical cyclone tracks detected from the ERA5 atmospheric analysis using version 13.2 of the CEOS/NSIDC extratropical cyclone tracking algorithm (as described in Crawford et al., 2021; https://doi.org/10.1175/MWR-D-20-0417.1). Storms are detected with a 3-h temporal resolution and 25-km spatial resolution for the period 1980-1989. All storms whose ending time (cyclolysis) occurs in the same month are grouped into a single CSV file and can be identified by their unique system number (sid). The monthly files are organized within folders for each year. See accompanying metadata file for more details (e.g., descriptions and units for each column). This algorithm explicitly tracks each center of a multi-center cyclones, but in this database, each multi-center cyclone is represented only by its primary center -- i.e., the entire storm system is being represented by the location and area. Only storm systems that meet the following four criteria are included: lifespan of at least 24 hours, track length of at least 1000 km, observed at least once over an elevation less than 500 m, and observed at least once at a distance of at least 500 km from its origin point are included.
Format	ZIP
Resource Category	data
URL	https://canwin-datahub.ad.umanitoba.ca/data/dataset/4be4d01a-a14b-483f-a1a6-6ead0974fa57/resource/83f7269e-c4f5-4690-bd50-71e4d0476890/download/cnect-nh-era5-1990-1999.zip
Name	Northern Hemisphere Extratropical Cyclone Tracks from ERA5: 1990-1999
Description	Folder containing CSV files that describe extratropical cyclone tracks detected from the ERA5 atmospheric analysis using version 13.2 of the CEOS/NSIDC extratropical cyclone tracking algorithm (as described in Crawford et al., 2021; https://doi.org/10.1175/MWR-D-20-0417.1). Storms are detected with a 3-h temporal resolution and 25-km spatial resolution for the period 1990-1999. All storms whose ending time (cyclolysis) occurs in the same month are grouped into a single CSV file and can be identified by their unique system number (sid). The monthly files are organized within folders for each year. See accompanying metadata file for more details (e.g., descriptions and units for each column). This algorithm explicitly tracks each center of a multi-center cyclones, but in this database, each multi-center cyclone is represented only by its primary center -- i.e., the entire storm system is being represented by the location and area. Only storm systems that meet the following four criteria are included: lifespan of at least 24 hours, track length of at least 1000 km, observed at least once over an elevation less than 500 m, and observed at least once at a distance of at least 500 km from its origin point are included.
Format	ZIP
Resource Category	data
URL	https://canwin-datahub.ad.umanitoba.ca/data/dataset/4be4d01a-a14b-483f-a1a6-6ead0974fa57/resource/8d5d2c9e-196b-4dbe-ac87-6a6cebcf0ae2/download/cnect-nh-era5-2000-2009.zip
Name	Northern Hemisphere Extratropical Cyclone Tracks from ERA5: 2000-2009

Description	Folder containing CSV files that describe extratropical cyclone tracks detected from the ERA5 atmospheric analysis using version 13.2 of the CEOS/NSIDC extratropical cyclone tracking algorithm (as described in Crawford et al., 2021; https://doi.org/10.1175/MWR-D-20-0417.1). Storms are detected with a 3-h temporal resolution and 25-km spatial resolution for the period 2000-2009. All storms whose ending time (cyclolysis) occurs in the same month are grouped into a single CSV file and can be identified by their unique system number (sid). The monthly files are organized within folders for each year. See accompanying metadata file for more details (e.g., descriptions and units for each column). This algorithm explicitly tracks each center of a multi-center cyclones, but in this database, each multi-center cyclone is represented only by its primary center -- i.e., the entire storm system is being represented by the location and area. Only storm systems that meet the following four criteria are included: lifespan of at least 24 hours, track length of at least 1000 km, observed at least once over an elevation less than 500 m, and observed at least once at a distance of at least 500 km from its origin point are included.
Format	ZIP
Resource Category	data
URL	https://canwin-datahub.ad.umanitoba.ca/data/dataset/4be4d01a-a14b-483f-a1a6-6ead0974fa57/resource/70bf8f2c-83da-4c88-bfbc-fe0a17a73796/download/cnect-nh-era5-2010-2019.zip
Name	Northern Hemisphere Extratropical Cyclone Tracks from ERA5: 2010-2019
Description	Folder containing CSV files that describe extratropical cyclone tracks detected from the ERA5 atmospheric analysis using version 13.2 of the CEOS/NSIDC extratropical cyclone tracking algorithm (as described in Crawford et al., 2021; https://doi.org/10.1175/MWR-D-20-0417.1). Storms are detected with a 3-h temporal resolution and 25-km spatial resolution for the period 2010-2019. All storms whose ending time (cyclolysis) occurs in the same month are grouped into a single CSV file and can be identified by their unique system number (sid). The monthly files are organized within folders for each year. See accompanying metadata file for more details (e.g., descriptions and units for each column). This algorithm explicitly tracks each center of a multi-center cyclones, but in this database, each multi-center cyclone is represented only by its primary center -- i.e., the entire storm system is being represented by the location and area. Only storm systems that meet the following four criteria are included: lifespan of at least 24 hours, track length of at least 1000 km, observed at least once over an elevation less than 500 m, and observed at least once at a distance of at least 500 km from its origin point are included.
Format	ZIP
Resource Category	data
URL	https://canwin-datahub.ad.umanitoba.ca/data/dataset/4be4d01a-a14b-483f-a1a6-6ead0974fa57/resource/ced01a53-f2ca-48fd-809c-cd99b83b7431/download/cnect-nh-era5-2020.zip
Name	Northern Hemisphere Extratropical Cyclone Tracks from ERA5: 2020

Description	Folder containing CSV files that describe extratropical cyclone tracks detected from the ERA5 atmospheric analysis using version 13.2 of the CEOS/NSIDC extratropical cyclone tracking algorithm (as described in Crawford et al., 2021; https://doi.org/10.1175/MWR-D-20-0417.1). Storms are detected with a 3-h temporal resolution and 25-km spatial resolution for the year 2020. All storms whose ending time (cyclolysis) occurs in the same month are grouped into a single CSV file and can be identified by their unique system number (sid). The monthly files are organized within folders for each year. See accompanying metadata file for more details (e.g., descriptions and units for each column). This algorithm explicitly tracks each center of a multi-center cyclones, but in this database, each multi-center cyclone is represented only by its primary center – i.e., the entire storm system is being represented by the location and area. Only storm systems that meet the following four criteria are included: lifespan of at least 24 hours, track length of at least 1000 km, observed at least once over an elevation less than 500 m, and observed at least once at a distance of at least 500 km from its origin point are included.
Format	ZIP
Resource Category	data
URL	https://canwin-datahub.ad.umanitoba.ca/data/dataset/4be4d01a-a14b-483f-a1a6-6ead0974fa57/resource/9b63b1ad-f5f1-4b4a-9c4b-8ca0a913a6b1/download/cnect-nh-era5-2021.zip
Name	Northern Hemisphere Extratropical Cyclone Tracks from ERA5: 2021
Description	Folder containing CSV files that describe extratropical cyclone tracks detected from the ERA5 atmospheric analysis using version 13.2 of the CEOS/NSIDC extratropical cyclone tracking algorithm (as described in Crawford et al., 2021; https://doi.org/10.1175/MWR-D-20-0417.1). Storms are detected with a 3-h temporal resolution and 25-km spatial resolution for the year 2021. All storms whose ending time (cyclolysis) occurs in the same month are grouped into a single CSV file and can be identified by their unique system number (sid). The monthly files are organized within folders for each year. See accompanying metadata file for more details (e.g., descriptions and units for each column). This algorithm explicitly tracks each center of a multi-center cyclones, but in this database, each multi-center cyclone is represented only by its primary center – i.e., the entire storm system is being represented by the location and area. Only storm systems that meet the following four criteria are included: lifespan of at least 24 hours, track length of at least 1000 km, observed at least once over an elevation less than 500 m, and observed at least once at a distance of at least 500 km from its origin point are included.
Format	ZIP
Resource Category	data
URL	https://canwin-datahub.ad.umanitoba.ca/data/dataset/4be4d01a-a14b-483f-a1a6-6ead0974fa57/resource/e9ef2608-e3f1-4edf-b987-281bd13b1e62/download/cnect-nh-era5-2022.zip
Name	Northern Hemisphere Extratropical Cyclone Tracks from ERA5: 2022

Description	Folder containing CSV files that describe extratropical cyclone tracks detected from the ERA5 atmospheric analysis using version 13.2 of the CEOS/NSIDC extratropical cyclone tracking algorithm (as described in Crawford et al., 2021; https://doi.org/10.1175/MWR-D-20-0417.1). Storms are detected with a 3-h temporal resolution and 25-km spatial resolution for the year 2022. All storms whose ending time (cyclolysis) occurs in the same month are grouped into a single CSV file and can be identified by their unique system number (sid). The monthly files are organized within folders for each year. See accompanying metadata file for more details (e.g., descriptions and units for each column). This algorithm explicitly tracks each center of a multi-center cyclones, but in this database, each multi-center cyclone is represented only by its primary center – i.e., the entire storm system is being represented by the location and area. Only storm systems that meet the following four criteria are included: lifespan of at least 24 hours, track length of at least 1000 km, observed at least once over an elevation less than 500 m, and observed at least once at a distance of at least 500 km from its origin point are included.
Format	ZIP
Resource Category	data
URL	https://canwin-datahub.ad.umanitoba.ca/data/dataset/4be4d01a-a14b-483f-a1a6-6ead0974fa57/resource/8bc6e96f-b50c-4fef-906c-16fc1ca98522/download/cnect-nh-era5-2023.zip
Name	Northern Hemisphere Extratropical Cyclone Tracks from ERA5: 2023
Description	Folder containing CSV files that describe extratropical cyclone tracks detected from the ERA5 atmospheric analysis using version 13.2 of the CEOS/NSIDC extratropical cyclone tracking algorithm (as described in Crawford et al., 2021; https://doi.org/10.1175/MWR-D-20-0417.1). Storms are detected with a 3-h temporal resolution and 25-km spatial resolution for the year 2023. All storms whose ending time (cyclolysis) occurs in the same month are grouped into a single CSV file and can be identified by their unique system number (sid). The monthly files are organized within folders for each year. See accompanying metadata file for more details (e.g., descriptions and units for each column). This algorithm explicitly tracks each center of a multi-center cyclones, but in this database, each multi-center cyclone is represented only by its primary center – i.e., the entire storm system is being represented by the location and area. Only storm systems that meet the following four criteria are included: lifespan of at least 24 hours, track length of at least 1000 km, observed at least once over an elevation less than 500 m, and observed at least once at a distance of at least 500 km from its origin point are included.
Format	ZIP
Resource Category	data
URL	https://canwin-datahub.ad.umanitoba.ca/data/dataset/4be4d01a-a14b-483f-a1a6-6ead0974fa57/resource/a5befcfe-1b51-48a4-ae05-159b8db5ef2c/download/cnect-nh-era5-2024.zip
Name	Northern Hemisphere Extratropical Cyclone Tracks from ERA5: 2024

Description	Folder containing CSV files that describe extratropical cyclone tracks detected from the ERA5 atmospheric analysis using version 13.2 of the CEOS/NSIDC extratropical cyclone tracking algorithm (as described in Crawford et al., 2021; https://doi.org/10.1175/MWR-D-20-0417.1). Storms are detected with a 3-h temporal resolution and 25-km spatial resolution for the year 2024. All storms whose ending time (cyclolysis) occurs in the same month are grouped into a single CSV file and can be identified by their unique system number (sid). The monthly files are organized within folders for each year. See accompanying metadata file for more details (e.g., descriptions and units for each column). This algorithm explicitly tracks each center of a multi-center cyclones, but in this database, each multi-center cyclone is represented only by its primary center – i.e., the entire storm system is being represented by the location and area. Only storm systems that meet the following four criteria are included: lifespan of at least 24 hours, track length of at least 1000 km, observed at least once over an elevation less than 500 m, and observed at least once at a distance of at least 500 km from its origin point are included.
Format	ZIP
Resource Category	data

Related Publications

Title	The Response of extratropical cyclone propagation in the Northern Hemisphere to global warming
URL	https://canwin-datahub.ad.umanitoba.ca/data/en/publication/the-response-of-extratropical-cyclone-propagation-in-the-northern-hemisphere-to-global-warming